

AD-A135 508

H/D EXCHANGE IN THE REACTION OF D2 WITH  
BIS(TRIPHENYLPHOSPHITE) (ACETYLAC...U) ROCHESTER UNIV  
NY DEPT OF CHEMISTRY B C WHITMORE ET AL. 04 NOV 83  
UROCHESTER/DC/83/TR-5 N00014-83-K-0154 F/G 7/2


1/1

UNCLASSIFIED

NL



END  
DATE  
1 54  
DTI



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

12

AD-A185-528

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0154

Task No. NR 634-742

TECHNICAL REPORT NO. 5

H/D Exchange in the Reaction of  $D_2$  with Bis(triphenylphosphite)  
(acetylacetonato)rhodium(I),  $Rh(P(OPh)_3)_2(acac)$

by

Bryan C. Whitmore and Richard Eisenberg

Prepared for Publication

in the

Journal of the American Chemical Society

University of Rochester

Department of Chemistry

Rochester, New York 14627

November 4, 1983

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

This document has been approved for public release  
and sale; its distribution is unlimited

DTIC  
ELECTE  
S DEC 7 1983 D

DTIC FILE COPY

B

88 12 07 056

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER <b>A1351308</b>	2. SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)	3. REPORT TYPE AND PERIOD COVERED Interim Technical Report	4. PERFORMING ORG. REPORT NUMBER
5. AUTHOR Bryan C. Whitmore and Richard Eisenberg	6. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Rochester Rochester, New York 14627	7. AUTHORING OR MONITORING NUMBER N00014-83-K-0154	8. PERFORMING ORG. NAME AND ADDRESS Navy Research Laboratory, VMC NR 634-742
9. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217	10. SECURITY CLASSIFICATION OF THIS REPORT Unclassified	11. SECURITY CLASSIFICATION OF THIS REPORT Unclassified	12. SECURITY CLASSIFICATION OF THIS REPORT Unclassified
13. ABSTRACT (Maximum 200 words) This document has been approved for public release and sale; its distribution is unlimited.			
14. DISTRIBUTION STATEMENT (For the abstract entered in block 13, if different from above) Prepared for publication in Journal of Am. Chem. Soc., in press			
15. SUPPLEMENTARY NOTES			
16. SECURITY CLASSIFICATION OF ABSTRACT (When Data Entered) Phosphite complexes, oxidative addition, acetylacetonate, NMR, metal hydrides, H/D exchange			
17. SECURITY CLASSIFICATION OF ABSTRACT (When Data Entered) The reaction of $\text{Rh}(\text{P}(\text{OPh})_2)_2(\text{acac})$ (I) with $\text{D}_2$ in benzene has been studied by $^1\text{H}$ NMR spectroscopy, and complex I has been found to undergo H/D exchange at the ortho positions of the coordinated phosphite ligands and at the central methine position of the acetylacetonate ligand. At $79^\circ\text{C}$ , the exchange reaction proceeds with the extent of deuterium incorporation into $\text{P}(\text{OPh})_2$ being the same as that into acac at all stages of the H/D exchange process. At $60^\circ\text{C}$ , deuterium incorporation into $\text{P}(\text{OPh})_2$ is initially more rapid than that seen into the acac ligand. The initial rate of deuterium incorporation into			

DD FORM 101 10/79 PREVIOUS EDITIONS ARE OBSOLETE  
GPO 1979-1-7-010-0001

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)  
 $\text{P}(\text{OPh})_2$  being the same as that into acac at all stages of the H/D exchange process. At  $60^\circ\text{C}$ , deuterium incorporation into  $\text{P}(\text{OPh})_2$  is initially more rapid than that seen into the acac ligand. The initial rate of deuterium incorporation into  $\text{P}(\text{OPh})_2$  by I in  $d_6$ -benzene under  $\text{D}_2$  at  $60^\circ\text{C}$  proceeds with a first order rate constant of  $9.6 \times 10^{-5} \text{ sec}^{-1}$ . A mechanism for this exchange process is proposed.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Contribution from the Department of Chemistry  
University of Rochester, Rochester, New York 14627

H/D Exchange in the Reaction of  $D_2$  with Bis(triarylphosphite)  
(acetylacetonate)rhodium(I),  $Rh(P(OR)_3)_2(acac)$

Bryan C. Whitmore and Richard Eisenberg\*


**ABSTRACT**

The reaction of  $Rh(P(OR)_3)_2(acac)$  (1) with  $D_2$  in benzene has been studied by  $^1H$  NMR spectroscopy, and complex 1 has been found to undergo H/D exchange at the ortho positions of the coordinated phosphite ligands and at the central methine position of the acetylacetonate ligand. At  $70^\circ C$ , the exchange reaction proceeds with the extent of deuterium incorporation into  $P(OR)_3$  being the same as that into acac at all stages of the H/D exchange process. At  $60^\circ C$ , deuterium incorporation into  $P(OR)_3$  is initially more rapid than that seen into the acac ligand. The initial rate of deuterium incorporation into  $P(OR)_3$  by 1 in  $d_6$ -benzene under  $D_2$  at  $60^\circ C$  proceeds with a first order rate constant of  $9.6 \times 10^{-4} \text{ sec}^{-1}$ . A mechanism for this exchange process is proposed. ←

10 to the minus 5<sup>th</sup> per sec

**Introduction**

Rhodium(I) complexes containing a single acetylacetonate (acac) ligand of the type  $RhLL'(acac)$ , where  $L, L' = CO, PPh_3, C_6H_5$ , were first reported in 1964.<sup>1</sup> Since that time numerous complexes containing different ligands, L, have been reported and studied. Many of these complexes are used as catalyst precursors for industrially important reactions such as olefin hydrogenation<sup>2</sup> and hydroformylation<sup>3,4</sup>, and have appeared in numerous patents.<sup>5</sup> One acac complex recently reported to catalyze arane hydrogenation is the phosphite complex,  $Rh(P(OR)_3)_2(acac)$ , 1.<sup>6</sup> Related rhodium(I) complexes containing phosphite ligands such as  $Rh(P(OR)_3)_2Cl$  and  $Rh(P(OR)_3)_3$  have been studied by Sarofield and Parrish,<sup>7</sup> and have been found to undergo H/D exchange under  $D_2$  at the ortho positions of the coordinated triaryl phosphite ligands.<sup>7</sup> This exchange reaction has been proposed by Parrish to proceed through an ortho-metalated intermediate 2 as shown in equation (1). Based on the reactivity of known rhodium(I) phosphite



complexes and the utilization of  $Rh(I)$ -acac complexes as homogeneous catalyst precursors, we have examined the reaction of complex 1,  $Rh(P(OR)_3)_2(acac)$ , with  $H_2$  and  $D_2$ . The results obtained from these reactions may aid in the elucidation of mechanisms for reactions catalyzed by complex 1 and other rhodium(I) acetylacetonate complexes.

### Experimental

$\text{Rh}(\text{P}(\text{OPh})_3)_2(\text{acac})$  was prepared by the literature method.<sup>8</sup> All experiments were performed in sealed NMR tubes in  $d_6$ -benzene.  $^1\text{H}$  NMR spectra were recorded on a Bruker WM-400 400 MHz instrument with chemical shifts reported in ppm relative to TMS.

**General procedure for  $^1\text{H}$  NMR experiments.** The rhodium complex,  $\text{Rh}(\text{P}(\text{OPh})_3)_2(\text{acac})$ , is placed in a 5 mm NMR tube which is evacuated on a high vacuum line. The tube is cooled to  $-196^\circ\text{C}$ . Liquid reactants (acetylacetone,  $\text{P}(\text{OPh})_3$ ) and  $d_6$ -benzene (0.5ml) are condensed into the NMR tube and the sample is then flame sealed under an  $\text{H}_2$  or  $\text{D}_2$  atmosphere. The reaction solution is heated at the indicated temperature and monitored by  $^1\text{H}$  NMR spectroscopy. Specific reaction conditions are summarized in Table I.

**Phosphite exchange reaction.** The NMR tube from reaction 1 (Table I) is broken open in air at the completion of the exchange reaction (98% deuterium incorporation). Half of the  $d_6$ -benzene solution of complex  $d_{13}\text{-1}$  (2.7mmol) is treated with  $\text{P}(\text{OPh})_3$  (1.5ml, 5.7mmol) and its  $^1\text{H}$  NMR spectrum is recorded immediately. The deuterium content of the ortho position of coordinated  $\text{P}(\text{OPh})_3$  is 46% within 5 minutes of mixing as determined through comparison of its integrated intensity with the integrated areas of the meta and para positions. The resonance due to the central methine position of acac remains unchanged.

### Results and Discussion

**$\text{H/D}$  exchange at  $75^\circ\text{C}$ .** When a  $d_6$ -benzene solution of  $\text{Rh}(\text{P}(\text{OPh})_3)_2(\text{acac})$ , **1**, is heated at  $75^\circ\text{C}$  under  $\text{D}_2$  (495 torr) (Reaction 1, Table I) a gradual change is observed in the 6.8-7.5 ppm region of its  $^1\text{H}$  NMR spectrum as shown in Figure 1. The aromatic region in the initial spectrum, (a), contains three multiplets assignable to the coordinated triphenyl phosphite ligands. The ortho hydrogens appear at 7.46ppm, and are split into a doublet by the meta hydrogens ( $J_{\text{ortho-meta}} = 8.0\text{Hz}$ ). The meta hydrogens give rise to a doublet of doublets at 7.01 ppm, and the


para hydrogens appear as a triplet at 6.85 ppm ( $J_{\text{para-meta}} = 7.4\text{Hz}$ ). Within several hours of heating the complex under  $\text{D}_2$ , the meta hydrogen resonance becomes increasingly complicated and the ortho hydrogen resonance decreases in integrated intensity compared to the meta and para hydrogens (spectrum (b)). In spectrum (c) the resonance due to the ortho hydrogens is nearly gone, and the meta hydrogen resonance now appears as a doublet ( $J = 7.4\text{Hz}$ ). The para hydrogen resonance remains unchanged throughout this reaction. These spectral changes are consistent with deuterium substitution into the ortho positions of the coordinated triphenyl phosphite ligands, and are similar to spectral changes in previous reports of  $\text{H/D}$  exchange in related  $\text{Rh}(\text{I})$  triphenyl phosphite complexes.<sup>7,9</sup> However, as deuterium is incorporated into the phosphite ligand in  $\text{Rh}(\text{P}(\text{OPh})_3)_2(\text{acac})$ , there occurs a concomitant decrease in the integrated intensity of the central methine proton of the acetylacetonate ligand indicating deuterium substitution into this position as well. At any point during this  $\text{D}$  substitution, the percentage of protons remaining in the methine position of the acac ligand is the same as the percentage of hydrogens seen in the ortho positions of the coordinated  $\text{P}(\text{OPh})_3$  ligand (see Figure 2). The only resonances observed in this exchange reaction are those attributable to  $\text{Rh}(\text{P}(\text{OPh})_3)_2(\text{acac})$  and its partially deuterated homologs. When the reaction is carried out in  $d_6$ -benzene under  $\text{H}_2$ , no evidence of hydride formation is observed, nor is any evidence of free acetylacetone or  $\text{P}(\text{OPh})_3$  obtained. Since complex **1** is the only species seen in solution by  $^1\text{H}$  NMR spectroscopy, any intermediates in the exchange process are present in only very small concentrations. The fact that the rate of  $\text{D}$ -incorporation into acac and  $\text{P}(\text{OPh})_3$  is the same indicates that a species common to both exchange processes, or related by ancillary equilibria, exists. If the two exchange processes were completely independent, proceeding without a common intermediate or equilibrium-related species, then two distinct rates of  $\text{D}$ -incorporation - one for  $\text{P}(\text{OPh})_3$

and one for acac - would be observed. This is clearly not the case for  $\text{Rh}(\text{P}(\text{OPh})_2)_2(\text{acac})$ . In addition, the fact that the extent of deuteration at the methine position of acac is always the same as that at the ortho positions of  $\text{P}(\text{OPh})_2$  indicates that the rate of D-incorporation into the latter is greater than that into acac. If the reverse were true - i.e., if D-incorporation into acac were faster than the deuteration of  $\text{P}(\text{OPh})_2$  - then the methine proton resonance would decrease in intensity more rapidly than the o-proton resonance of  $\text{P}(\text{OPh})_2$ . This conclusion concerning relative rates of deuterium incorporation is supported by studies of exchange at 60°C described below.

In order to probe more fully the nature of the H/D exchange reactions at 75°C, we performed additional reactions in the presence of added  $\text{P}(\text{OPh})_2$  and/or acetylacetonate ligand. When complex 1 is placed under  $\text{D}_2$  in the presence of 1 equiv  $\text{P}(\text{OPh})_2$ , deuterium incorporation into both coordinated and free  $\text{P}(\text{OPh})_2$  is observed, as well as into the methine position of the acac ligand. A small amount of free acetylacetonate is also detected. Similarly, when 1 is heated at 75°C under  $\text{D}_2$  with 2.7 equiv of Hacac added, exchange into both coordinated and free acac is seen, as well as into the ortho positions of coordinated  $\text{P}(\text{OPh})_2$ . H/D exchange into free and coordinated ligands is also seen when the same reaction is run with 1 equiv  $\text{P}(\text{OPh})_2$  and 6 equiv Hacac added. Clearly, ligand exchange is a factor in analyzing the H/D exchange reactions in the  $\text{Rh}(\text{P}(\text{OPh})_2)_2(\text{acac})$  system.


The facility of phosphite exchange with 1 is demonstrated by the following reaction. When 2.1 equivalents of  $\text{P}(\text{OPh})_2$  are added to the  $d_5$ -benzene solution of  $d_{13}$ -1 (90%, formed at the completion of reaction 1, Table I) at room temperature there is an immediate reappearance in the  $^1\text{H}$  NMR spectrum of the resonance due to the ortho protons of coordinated  $\text{P}(\text{OPh})_2$ . Within five minutes, the integrated intensity of the ortho protons from coordinated  $\text{P}(\text{OPh})_2$  indicates a

46% deuterium content which corresponds to complete scrambling of labelled and unlabelled  $\text{P}(\text{OPh})_2$  ligands (equation 2).





The facile exchange of free and coordinated phosphite ligands in this system contrasts with a recent study using  $^{31}\text{P}$  NMR spectroscopy which concludes that phosphite exchange for 1 does not occur or occurs only very slowly. While this ligand exchange may be too slow to be observed by room temperature NMR line broadening, it is readily observed at room temperature using labelled phosphite ligands. The results of this  $\text{P}(\text{OPh})_2$  exchange experiment clearly show that 1 exchanges  $\text{P}(\text{OPh})_2$  more rapidly than 1 undergoes deuterium incorporation reactions. When the reaction mixture (1 +  $\text{D}_2$ ) is maintained at room temperature (25°C) there is no change in its  $^1\text{H}$  NMR spectrum over the course of several hours. Deuterium incorporation into the coordinated ligands is detected only when the benzene solution of 1 +  $\text{D}_2$  is heated.

H/D exchange at 60°C. When the reaction of  $\text{Rh}(\text{P}(\text{OPh})_2)_2(\text{acac})$  with  $\text{D}_2$  is carried out at 60°C, changes are seen in its  $^1\text{H}$  NMR spectrum similar to those observed at 75°C. However, at this lower temperature, D-incorporation into the phosphite ligand is initially much more rapid than substitution into the acac ligand as illustrated in Table II. Thus, deuterium substitution occurs primarily on the  $\text{P}(\text{OPh})_2$  ligand during the early stages of the exchange reaction at 60°C. This reaction, shown as equation 3, can be considered a simple isotope exchange



reaction, which is the first of a series of H/D exchanges of the general form  $d_n-1 + D_2 \rightarrow d_{n+1}-1 + HD$ , where  $n=0-12$ . However, as the exchange process continues, the kinetics of H/D exchange grows in complexity due to the increasing probability of back-reactions and degenerate exchanges as shown in eqs (4)-(9). After several exchanges have occurred there



is an increase first in the concentration of HD and subsequently in the concentration of  $H_2$ . Indeed, during the exchange reactions both HD and  $H_2$  are detected by  $^1H$  NMR spectroscopy. A 1:1:1 triplet ( $J = 42$  Hz) at 4.42 ppm is seen for HD, while  $H_2$  gives rise to a singlet at 4.45 ppm (in  $d_6$ -benzene). As the concentrations of these molecules increase, there is an increase in the likelihood that they will become involved in the exchange processes (4) - (9). Although  $D_2$  is present in the excess in the gas phase above the reaction solutions, its concentration in the benzene solutions is nearly the same as that of the rhodium complex,  $\lambda$ . For example, in reaction 5 (Table I) under 517 torr pressure the concentration of  $D_2$  in solution is roughly 2M while the concentration of  $\lambda$  is 4.8M. Due to the low surface area of benzene in the NMR tube, as  $D_2$  is consumed in solution, it may

not be immediately replaced by  $D_2$  from the gas phase and the relative concentrations of HD and  $H_2$  become larger than expected. Thus as the reaction proceeds, it becomes mass transfer limited.

Despite the obvious complexity of this exchange process, an estimate of the rate of H/D exchange in coordinated  $P(OPh)_3$  during the early stages of the reaction  $\lambda + D_2$  can be obtained using eqn (10) derived by Calvin<sup>10</sup> and other workers<sup>11</sup> which deals with the rate of single isotope exchange.

$$\ln \frac{[A_0]}{[A]} = \frac{1}{a+b} (a+b)t = kt \quad (10)$$

$\lambda$  and  $\bar{\lambda}$  are defined in equation (3)

$[A_0]$  = concentration of A at  $t=0$

$a = [D_2] + [HD]$

$b = [A] + [B]$


A plot of  $\ln ([A_0]/[A])$  vs  $t$  for reaction 5, shown in Figure 3, is linear during the initial stages of the reaction, and yields a value for  $k$  in eqn (10) of  $9.6 \times 10^{-4} \text{ sec}^{-1}$ .

#### Mechanism of H/D Exchange

Any mechanism drawn for the H/D exchange reactions of complex  $\lambda$  must account for the results obtained in the experiments described above which are summarized as follows: (1) there is a facile room temperature exchange of free and coordinated triphenyl phosphite; (2) deuterium is incorporated into both  $P(OPh)_3$  and acetylcetonate ligands; (3) deuterium is incorporated into both free and coordinated ligands; (4) the relative rates of D-substitution into  $P(OPh)_3$  and acac are the same at 70°C.; (5) at 60°C, the rate of deuteriation of  $P(OPh)_3$  is faster than that of acac; (6) during the reaction of  $\lambda$  with  $D_2$  or  $H_2$ , no species except for  $\lambda$  is detected by  $^1H$  NMR spectroscopy; and (7) during the reaction of  $\lambda$  with  $D_2$  in the presence of excess  $P(OPh)_3$ , free acetylcetonate is observed by  $^1H$  NMR.




The incorporation of deuterium into the ortho-position of coordinated  $P(OPh)_3$  must involve the activation of both the  $D_2$  molecule and the ortho-phenyl position of the phosphite ligand. Both of these processes occur most logically via oxidative additions to  $Rh(I)$  centers. These oxidative additions have been proposed by Parshall to explain H/D exchange in similar rhodium phosphite complexes.<sup>7</sup> The incorporation of deuterium into coordinated acac on the other hand requires the formation of acetylacetone, either free or weakly coordinated, via the reductive elimination of Rhac from a Rh-B species as shown in equation (III).



If the oxidative addition of  $D_2$  to complex 1 were to occur first, then one forms species 2. The reductive elimination of  $BDac$  from 2 yields species 3 which can then undergo reversible orthometallation, thereby incorporating deuterium into the phosphite ligand. In this sequence,  $1 \rightleftharpoons 2 \rightleftharpoons 3$ , one would expect the incorporation of deuterium into  $acac$  to be observed immediately, and the rate of this exchange to be as fast as, if not faster than, the rate of  $\beta$ -substitution into the phosphite ligand. However, this situation is not observed. The experimental results indicate that the rate of substitution of deuterium into  $acac$  is


clear than the incorporation of deuterium into  $P(OH)_3$ . Therefore, the first step of the H/D exchange process must be the arthmetallation of the triphenyl phosphite ligand,  $\underline{1} \rightarrow \underline{2}$ . The reductive elimination of Haccac from the rhodium(III) center leads to the regeneration of a Rh(I) species,  $\underline{2}$ . Deuterium is then incorporated into the triphenyl phosphite ligand via the oxidative addition of  $D_2$ ,  $\underline{2} \rightarrow \underline{3}$ , followed by reductive elimination of the aryl and deuterium ligands,  $\underline{3} \rightarrow \underline{4}$ , for deuterium incorporation into acac to reflect the extent of deuteration of the  $P(OH)_3$  ligand, the source of deuterium for the acetylacetonate ligand must be the ortho positions of the phosphite ligand. This may be provided by rapid and reversible interconversions between  $\underline{3}$  and  $\underline{4}$  in which the deuteride of  $\underline{3}$  is scrambled among all of the ortho positions of  $P(OH)_3$ , or it may be provided by the conversion of  $\underline{1}$  to  $\underline{2}$ . In any case, at 75°C there is no observable difference in the extent of deuteration of the two ligands, and the cycle of Scheme 1 is traversed smoothly. At 60°C, however, D-incorporation into acac proceeds more slowly than into  $P(OH)_3$ , suggesting that interconversions of  $\underline{3} \rightleftharpoons \underline{4} \rightarrow \underline{2}$  occur more readily than the readdition of Haccac to the rhodium center either  $\underline{3} \rightarrow \underline{4}$  or  $\underline{2} \rightarrow \underline{2}$ . This readdition is required in order to obtain deuterium incorporation into the methine position of the acetylacetonate ligand via  $\underline{4} \rightarrow \underline{5}$  or  $\underline{4} \rightarrow \underline{2} \rightarrow \underline{5}$ .

In the H<sub>2</sub>O exchange acetylacetonate may remain weakly bound to the rhodium center as suggested in Scheme I or it may dissociate entirely from the metal center. If Hacac were to remain weakly coordinated, then it would be expected to adapt coordination mode I or II as shown below. If Hacac were totally dissociated from Rh, then it would be expected to bind to the metal center via I or II prior to the



formation of the acac chelate ring. For D-incorporation into acac, there must occur the formation of either weakly coordinated or free Dacac which can achieve H/D exchange via keto-enol tautomerism. The observation of a small amount of free acetylacetone by  $^1\text{H}$  NMR spectroscopy when  $\text{P(OPh)}_3$  is added to the reaction solution (reaction 2 of Table I) shows that complete dissociation of Hacac is occurring to a limited extent.

The observation that Hacac can be reductively eliminated from the rhodium center is very important because of the interest in rhodium acetylacetonate complexes as homogeneous catalyst precursors. The loss of the acac ligand from rhodium has been observed previously but under more severe conditions (e.g., 40 atm 1:1:1  $\text{CO}/\text{H}_2/\text{C}_2\text{H}_4$  for  $\text{Rh}(\text{CO})_2(\text{acac})^3$ ). We have shown that the reductive elimination of Hacac (or Dacac) from  $\text{Rh}(\text{P(OPh)}_3)_2(\text{acac})$  occurs under relatively mild conditions (60°C, 360 torr  $\text{H}_2$ ). In a metal acac complex, the acac ligand is a formal 3e-donor, therefore the reductive elimination of Hacac and subsequent dissociation from the metal center reduces the electron count of the metal center by 4. The loss of this bidentate ligand via reductive elimination and dissociation therefore generates a high degree of coordinative unsaturation at the rhodium center. The ease of this ligand displacement may explain the usefulness of complexes of this type as catalyst precursors. The dissociation of Hacac from species 3 in Scheme I generates a highly unsaturated three-coordinate  $\text{Rh(I)}$  complex 2. A complex of this type may be involved in the hydrogenation of aromatics recently reported to be catalyzed by complex 2 under 10 atm of  $\text{H}_2$  at 80°C. Similar rhodium phosphite complexes of the type  $(\text{H Rh}(\text{P}(\text{O}-i\text{-Pr})_3)_2)_2$  are known to be exceptionally active catalyst precursors for olefin hydrogenation reactions.<sup>12</sup> The rhodium tri(isopropyl)phosphite dimer is closely related to species 2 in Scheme I by the proposed sequence of reactions shown in equation 12, where  $\text{L} = \text{P(OPh)}_3$ .



Thus, the catalytic activity of analogous rhodium acac complexes is most likely a result of the ease of acac displacement via Hacac reductive elimination and dissociation. A closer examination of other catalytic reactions involving  $\text{Rh}(\text{acac})$  complexes as catalyst precursors will undoubtedly support this conclusion.

**Acknowledgment.** We wish to thank the Office of Naval Research for support of this work and Johnson-Matthey for a generous loan of precious metal salts. BCN acknowledges a Sherman Clarke Fellowship.

## References

1. Bonati, F.; Wilkinson, G. *J. Chem. Soc.* **1964**, 3156.
2. Shobaldova, A. B.; Bystronina, V. I.; Krutsova, V. N.; Khidol, N. L. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1975**, 2101.
3. King, R. D.; King, A. D. Jr.; Iqbal, N. Z. *J. Am. Chem. Soc.* **1979**, *101*, 4093.
4. Hartley, F. R.; Murray, S. G.; Nicholson, P. H. *J. Mol. Cat.* **1982**, *16*, 363.
5. a) Kaplan, L.; Pruett, R. L. (Union Carbide Corp.) *U.S. 4,199,621*, 22 Apr. **1980**.  
b) Kaplan, L. (Union Carbide Corp.) *U.S. Appl. 805, 993*, 30 June **1978**.  
c) Kaplan, L.; Cotton, F. A. (Union Carbide Corp.) *U.S. 4,188,330*, 12 Feb. **1980**.
6. Pieta, D.; Trzeciak, A. M.; Zidzinski, J. J. *J. Mol. Cat.* **1983**, *18*, 193.
7. Barefield, E. K.; Parrish, G. W. *Inorg. Chem.* **1972**, *11*, 964.
8. Trzeciak, A. M.; Zidzinski, J. J. *Inorg. Chim. Acta, Lett.* **1982**, *64*, L267.
9. Parrish, G. W.; Knott, M. H.; Schum, R. A. *J. Am. Chem. Soc.* **1963**, *91*, 4990.
10. Buffield, R. B.; Calvin, M. J. *J. Am. Chem. Soc.* **1966**, *88*, 947.
11. Moore, J. M.; Pearson, R. G. "Kinetics and Mechanism," John Wiley and Sons; New York, **1961**, p. 311 and references therein.
12. Sivak, A. J.; Huttert, E. L. *J. Am. Chem. Soc.* **1979**, *101*, 4670.

Table 1. Conditions for H/D exchange reactions

Reaction	T (°C)	Reactants (μmol)			
		Complex 1	Hexac	P(OPh) <sub>3</sub>	D <sub>2</sub>
1	75	5.4	-	-	53 <sup>a</sup>
2	75	6.0	-	6.0	52
3	75	3.6	9.8	-	53
4	75	6.0	3.0	6.0	51
5	60	2.4	-	-	56
6	60	3.0	-	-	38

<sup>a</sup> This corresponds to a D<sub>2</sub> pressure of 495 torr at 298°K

Table II. S-butarium incorporation into coordinated ligands at 60°C.

Reaction <sup>a</sup>	Time (hr)	S - B		$k^c$ (sec <sup>-1</sup> )
		AGAC	P(OPh) <sub>3</sub>	
5	0.5	3	14	$9.6 \times 10^{-5}$
	1.4	3	38	
	3.0	5	46	
	6.6	7	52	
	20 <sup>b</sup>	9	52	
6	0.3	3	6	$6.7 \times 10^{-5}$
	1.1	3	21	
	4.1	5	28	
	5.9	8	31	
	9.1	23	30	
	16.3	40	36	
	38.0	84	37	

<sup>a</sup> Reaction conditions given in Table I. <sup>b</sup> After 6.6 hr reaction at 60°, the NMR tube is maintained at room temperature for the following 13.2 hr. <sup>c</sup> Rate constant  $k$  from equation (10) corresponding to initial distribution.

Scheme I. Proposed mechanism for M/D exchange







Figure 1: Aromatic regions of the 400-MHz  $^1\text{H}$  NMR Spectra of Complex 1 under  $\text{D}_2$ .  
 $\ast = \text{C}_6\text{H}_5$ .

Figure 2: Plot of deuterium-incorporation into the  $\text{P(OPh)}_3$  and acetylacetonate ligands of complex 1 for reaction 1 (Table I.) vs. time.

Figure 3: Plot of  $\ln ([A_0]/[A])$  vs. time for reaction 5 (Table I.).





END

DATE  
FILMED

1 84

DTIC